

SU 1723125

L3 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2005 ACS on STN

AN 1993:75205 CAPLUS

DN 118:75205

Bis-N-aziridinealkanes as mutagens

IN Vasil'eva, S. V.; Kadorkina, G. K.; Kostyanovskii, R. G.; Makhova, E. V.; Pavlova-Rezakova, A. G.

PA Semenov, N. N., USSR

SO U.S.S.R.

From: Izobreteniya 1992, (12), 129.

DT Patent

LA Russian

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	SU 1723125	A1	19920330	SU 1989-4767001	
	19891208 <--				
PRAI	SU 1989-4767001		19891208		
AB	Bis-N-aziridinealkanes (I; n = 3-8, 12) are mutagens.				

IT Alkanes, compounds

RL: BIOL (Biological study)

(bisaziridine derivs., as mutagens)

IT Mutagens

(bisaziridinealkanes)

IT 18924-57-7 25781-25-3 40717-38-2 56522-41-9 134753-77-8

134753-78-9 134753-79-0

RL: BIOL (Biological study)

(as mutagen)

DIALOG(R)File 351:Derwent WPI

(c) 2005 Thomson Derwent. All rts. reserv.

009398212

WPI Acc No: 1993-091686/ 199311

Alpha, omega-bis-N-aziridino-alkane(s) used as chemical mutagens – are prepd. by reacting alpha, omega-diaminoalkane with dihalo-ethane in dichloroethane or benzene in presence of alkali

Patent Assignee: CHEM PHYS INST

Inventor: KADORKINA G K; KOSTYANOVSKII R G; VASILEVA S V

Number of Countries: 001 Number of Patents: 001

Patent Family:

Patent No	Kind	Date	Applicat No	Kind	Date	Week
SU 1723125	A1	19920330	SU 4767001	A	19891208	199311 B

Priority Applications (No Type Date): SU 4767001 A 19891208

Patent Details:

Patent No	Kind	Lan	Pg	Main IPC	Filing Notes
SU 1723125	A1		4	C12N-015/01	

Abstract (Basic): SU 1723125 A

Alpha, omega -bis-N- aziridinoalkanes of formula (I), where n = 3-8 and 12 is obt'd. by

SU 1723125 cont.

reacting alpha, omega-diaminoalkane with 1,2-dihaloethane in 1,2-dichloroethane or benzene, in the presence of 50% alkali, at 60-70 deg.C.

USE/ADVANTAGE - In genetics: (I) find novel use as chemical mutagens. They are safer and more active than the parent aziridine of formula (II), and are used as mutagens for bacterial cells of *Salmonella typhimurium* TA 100 with the genotype hisG46uvrBrfa pKM 101. The cpd. boils at much higher temp. of 190-220 deg.C, compared with (II) which boils at 50 deg.C.
Bul.12/30.3.92